

$B \rightarrow \pi\pi$ decays***T. N. Pham**

*Centre de Physique Théorique,
Centre National de la Recherche Scientifique, UMR 7644,
Ecole Polytechnique, 91128 Palaiseau Cedex, France*

ABSTRACT

The branching ratios and CP asymmetries of $B \rightarrow \pi\pi$ decays measured in B factory experiments indicate a large ratio of color-suppressed (C) to color-allowed (T) tree contributions and a large relative phase between the tree and penguin amplitude. In this talk, I would like to report on a recent analysis to see whether the large C/T ratio can be explained within the QCD based model with i) a large contribution from the annihilation terms or with ii) large final-state-interaction phase between two different isospin amplitudes. We show that the current experimental data do not exclude either possibility but we may be able to distinguish these two effects in future measurements of direct CP asymmetry in $B \rightarrow \pi^0\pi^0$ decay.

* Talk given at First Workshop On Theory, Phenomenology And Experiments In Heavy Flavor, Anacapri, Italy, May 29-31, 2006

1 Introduction

The large $B \rightarrow \pi^0\pi^0$ branching ratio measured at Babar and Belle indicates that the color-suppressed amplitude is large compared with predictions from the short-distance effective non-leptonic weak Hamiltonian. the ratio of the color-suppressed to color-allowed tree amplitude C/T is proportional to a_2/a_1 with $a_1 = C_1 + C_2/N_c$ and $a_2 = C_2 + C_1/N_c$ being the effective Wilson coefficients for the charged and neutral current tree-level color-singlet operator O_1 and O_2 respectively. At the renormalization scale $\mu \approx m_b[1]$, $C_1 \approx 1.10, C_2 \approx -0.250$ (at NLO) for non-leptonic B decay, a_2 therefore becomes much smaller than a_1 . Thus in the absence of higher order QCD radiative corrections and non-factorisable contributions, B decays with neutral particles in the final state are suppressed compared to decays with charged particles in the final state and are called color-suppressed decays. This rule is known to be violated in D decays and also in B decays with charmed meson in the final state. Since we expect QCD Factorization (QCDF)[2] to work better for charmless two-body B decays with energetic light hadrons in the final state, the large C/T ratio extracted from the BaBar and Belle $B \rightarrow \pi\pi$ measurements compared with theoretical predictions from factorization models is a surprise: The suppression of the $B^0 \rightarrow \pi^+\pi^-$ and the enhancement of the $B^0 \rightarrow \pi^0\pi^0$ decays. Also the large measured CP asymmetry in $B^0 \rightarrow \pi^+\pi^-$ implies a large relative phase between the tree and penguin amplitude and suggests that strong final state interaction effect could play an important role in $B \rightarrow \pi\pi$ decays which are governed not only by weak interaction but also by strong interactions. Therefore an understanding of the strong interaction effects in these processes is crucial for extracting the weak phase and ultimately possible new physics which could make additional contributions to the one-loop penguin amplitude [3]. In this talk, I would like to report on a recent work[4] in which we investigate two possible mechanisms for the enhancement of C/T in $B \rightarrow \pi\pi$ decays: i) the higher order correction of QCDF and ii) the effect of final state interaction (FSI) and show that in QCDF annihilation terms could play an important role in the enhancement of C/T but alternatively, a large FSI rescattering phase could also enhance C/T in QCDF with moderate annihilation contributions.

2 The decay amplitude from experiments

With the recent measurements of the branching ratios for the $B^0 \rightarrow \pi^+\pi^-$, $B^0 \rightarrow \pi^0\pi^0$ and $B^+ \rightarrow \pi^+\pi^0$ as well as the mixing-induced $S_{\pi^+\pi^-}$ and direct $C_{\pi^+\pi^-}$ CP asymmetries in $B^0 \rightarrow \pi^+\pi^-$ decays, it is now possible to determine the ratio of penguin to tree amplitude and the relative phase between tree and penguin amplitude as done in many previous model-independent analysis [5]-[18].

In the c-convention[5], the $B \rightarrow \pi\pi$ amplitudes are

$$A(B^0 \rightarrow \pi^+\pi^-) = T e^{i\delta_T} e^{i\gamma} + P e^{i\delta_P} \quad (1)$$

$$\sqrt{2}A(B^0 \rightarrow \pi^0\pi^0) = C e^{i\delta_C} e^{i\gamma} - P e^{i\delta_P} \quad (2)$$

$$\sqrt{2}A(B^+ \rightarrow \pi^+\pi^0) = (T e^{i\delta_T} + C e^{i\delta_C}) e^{i\gamma} \quad (3)$$

where $\delta_{T,C,P}$ are respectively the strong phase for the tree T , color-suppressed C and penguin P amplitude, γ being the CP violating weak phase. To determine C/T and the relative phase between C and T , we have 5 measured observables [19] (all branching ratios quoted are CP-

averaged)

$$S_{\pi^+\pi^-} = -0.50 \pm 0.12 \quad (4)$$

$$C_{\pi^+\pi^-} = -0.37 \pm 0.10 \quad (5)$$

$$\text{Br}(\pi^+\pi^-) = (4.5 \pm 0.4) \times 10^{-6} \quad (6)$$

$$\text{Br}(\pi^0\pi^0) = (1.45 \pm 0.29) \times 10^{-6} \quad (7)$$

$$\text{Br}(\pi^+\pi^0) = (5.5 \pm 0.6) \times 10^{-6} \quad (8)$$

where $S_{\pi^+\pi^-}$ and $C_{\pi^+\pi^-}$ are respectively the $B^0 - \bar{B}^0$ mixing-induced and direct CP asymmetries in the expression for the time-dependent CP asymmetry of $B \rightarrow \pi^+\pi^-$:

$$A_{\pi^+\pi^-}(t) = S_{\pi^+\pi^-} \sin(\Delta M_B t) - C_{\pi^+\pi^-} \cos(\Delta M_B t) \quad (9)$$

In terms of T , C and P , we have:

$$R S_{\pi^+\pi^-} = \sin 2\alpha + 2 \sin(\beta - \alpha) \cos \delta_{PT} \left(\frac{P}{T} \right) - \sin 2\beta \left(\frac{P}{T} \right)^2 \quad (10)$$

$$R C_{\pi^+\pi^-} = 2 \sin(\alpha + \beta) \sin \delta_{PT} \left(\frac{P}{T} \right) \quad (11)$$

where

$$R = 1 - 2 \cos(\alpha + \beta) \cos \delta_{PT} \left(\frac{P}{T} \right) + \left(\frac{P}{T} \right)^2 \quad (12)$$

The CP-averaged branching ratios are given by:

$$R_{00} = \frac{2\text{Br}(\pi^0\pi^0)}{\text{Br}(\pi^+\pi^-)} = \frac{1}{R} \left[\left(\frac{C}{T} \right)^2 + \left(\frac{P}{T} \right)^2 - 2 \cos(\delta_{PT} - \delta_{CT}) \cos \gamma \left(\frac{C}{T} \right) \left(\frac{P}{T} \right) \right] \quad (13)$$

$$R_{+-} = \frac{2\text{Br}(\pi^+\pi^0)\tau_{B^0}}{\text{Br}(\pi^+\pi^-)\tau_{B^+}} = \frac{1}{R} \left[1 + 2 \cos \delta_{CT} \left(\frac{C}{T} \right) + \left(\frac{C}{T} \right)^2 \right] \quad (14)$$

To have an idea of the size of the relative phase δ_{PT} , let us take as an approximation, $\alpha \approx 90^\circ$ and neglect the $(P/T)^2$ term in Eq.(10), we have

$$\tan \delta_{PT} \approx -C_{\pi^+\pi^-}/S_{\pi^+\pi^-} \quad (15)$$

which gives $\delta_{PT} = -36.5^\circ$ more or less in agreement with the more precise value -41.3° for $\gamma = 67^\circ$ obtained in Table 1. This simple calculation shows that there is a large relative penguin-tree strong phase in $B \rightarrow \pi\pi$ decays. Table 1 also shows the variation of P/T , R and δ_{PT} with γ and CP asymmetries within the uncertainties in the measured CP asymmetries and the determination of γ .

Figure 1 shows the allowed region for C/T and δ_{CT} plotted with the central values of the CP asymmetries measurements. Note that the overlap of the solid (R_{00}) and the dashed (R_{+-}) bounds shift towards the larger C/T region as γ becomes larger. R_{+-} allows relatively small value of C/T while R_{00} leads to a more strict constraint, $C/T \gtrsim 0.5$. The overlap region is distributed in a large range of δ_{CT} . Note also errors due to $(S_{\pi^+\pi^-}, C_{\pi^+\pi^-})$ are $\pm$ a few % in C/T and $\pm 20^\circ$ in δ_{CT} . This analysis shows clearly that C/T is large compared to the leading order QCD corrections ($C/T = 1/N_c$). In the next section we present results of QCDF calculation for C/T .

γ ($S_{\pi^+\pi^-}, C_{\pi^+\pi^-}$)	27°	37°	47°	57°	67°	77°	87°
(-0.62, -0.47)	0.53 -155° 0.43	0.38 -131° 0.75	0.36 -92.6° 1.10	0.46 -61.7° 1.45	0.63 -45.0° 1.75	0.81 -35.5° 1.96	0.98 -29.5° 2.06
(-0.62, -0.27)	0.49 -166° 0.40	0.29 -147° 0.69	0.19 -92.1° 1.03	0.32 -43.4° 1.35	0.52 -27.5° 1.63	0.72 -20.5° 1.82	0.91 -16.6° 1.91
(-0.50, -0.37)	0.55 -164° 0.35	0.38 -149° 0.62	0.26 -115° 0.92	0.29 -66.6° 1.21	0.44 -41.3° 1.46	0.62 -29.8° 1.63	0.80 -23.4° 1.72
(-0.38, -0.47)	0.61 -164° 0.33	0.45 -150° 0.58	0.33 -125° 0.85	0.31 -86.9° 1.12	0.41 -56.7° 1.35	0.56 -40.4° 1.51	0.72 -31.2° 1.58
(-0.38, -0.27)	0.59 -170° 0.31	0.40 -162° 0.55	0.23 -140° 0.81	0.17 -79.1° 1.07	0.31 -37.8° 1.28	0.49 -24.1° 1.44	0.67 -17.8° 1.51

Table 1: Determination of P/T (upper value), δ_{PT} (middle value) and R (bottom value) using experimental results for $S_{\pi^+\pi^-} = (-0.50 \pm 0.12)$ and $C_{\pi^+\pi^-} = (-0.37 \pm 0.10)$ for given values of γ , $\gamma = (27^\circ \sim 87^\circ)$.

3 C/T in QCD Factorization

Neglecting electroweak penguin contributions, T , C and P for $B \rightarrow \pi\pi$ decays in QCD Factorization [2, 20] are given by:

$$Te^{i\delta_T}e^{i\gamma} \propto \lambda_u^*(a_1 + b_1 + \hat{a}_4^u), \quad (16)$$

$$Ce^{i\delta_C}e^{i\gamma} \propto \lambda_u^*(a_2 - b_1 - \hat{a}_4^u), \quad (17)$$

$$Pe^{i\delta_P} \propto \lambda_c^*\hat{a}_4^c \quad (18)$$

$$\hat{a}_4^p = a_4^p + r_\chi a_6^p + 2b_4 + b_3 \quad (19)$$

$r_\chi = 2m_\pi^2/(2m_b m_q) \simeq 1.24$ and $m_q \equiv (m_u + m_d)/2$.

$$\lambda_u = V_{ub}V_{ud}^* \simeq A\lambda^3(\rho - i\eta) \quad (20)$$

$$\lambda_c = V_{cb}V_{cd}^* \simeq -A\lambda^3. \quad (21)$$

where a_i contain the factorisable term, the vertex correction, hard-scattering correction and penguin correction and b_i are annihilation contributions. Using the following parameters,

$$\begin{aligned} \mu &= 4.2 \text{ GeV}, \quad m_q(2\text{GeV}) = 0.0037 \text{ GeV}, \quad \lambda_B = 0.35 \text{ GeV}, \\ |\lambda_u/\lambda_c| &= |\rho + i\eta| = 0.09, \quad \alpha_2^\pi = 0.1 \end{aligned} \quad (22)$$

we find

$$a_1 = 1.02e^{i0.8^\circ} - 0.014\rho_H e^{i\phi_H} \quad (23)$$




Figure 1: Allowed region for δ_{CT} (x -axis) versus C/T (y -axis), from bounds for R_{00} and R_{+-} . Values for P/T and δ_{PT} are from the central value of $(S_{\pi^+\pi^-}, C_{\pi^+\pi^-}) = (-0.50, -0.37)$. The three solid lines represent $R_{00} = 0.64 - 0.14, 0.64, 0.64 + 0.14$, the three dashed lines represent $R_{+-} = 2.27 - 0.32, 2.27, 2.27 + 0.32$. The overlap of solid and dashed bounds are the allowed region for C/T and δ_{CT} .

$$a_2 = 0.21e^{-i23^\circ} + 0.081\rho_H e^{i\phi_H} \quad (24)$$

$$a_4^u + r_\chi a_6^u = -0.097e^{i21^\circ} + 0.0010\rho_H e^{i\phi_H} \quad (25)$$

$$a_4^c + r_\chi a_6^c = -0.10e^{i7^\circ} + 0.0010\rho_H e^{i\phi_H} \quad (26)$$

and

$$b_1 = 0.027 + 0.063\rho_A e^{i\phi_A} + 0.0085(\rho_A e^{i\phi_A})^2 \quad (27)$$

$$b_3 = -0.0067 - 0.021\rho_A e^{i\phi_A} - 0.015(\rho_A e^{i\phi_A})^2 \quad (28)$$

$$b_4 = -0.0019 - 0.0046\rho_A e^{i\phi_A} - 0.0006(\rho_A e^{i\phi_A})^2 \quad (29)$$

We have made a complete analysis of C/T covering all the parameter space of ρ 's and ϕ 's. In the limit of $\rho_{H,A} = 0$. We find

$$\frac{C_0}{T_0} e^{i\delta_{CT_0}} = 0.29e^{-i8.5^\circ} \quad (30)$$

$\rho_{H,A}$ and $\phi_{A,H}$ represent the divergent end-point singularity terms for the hard scattering function and the annihilation diagrams which introduce large theoretical uncertainties in the QCD calculation. Fig.2 shows δ_{CT} versus C/T for $-\pi < \phi_{A,H} < \pi$ with fixed ρ_H and ρ_A which would vary, say, in the ranges of $|\rho_{A,H}| < 1 \sim 2$. We see that quite a large range of C/T and δ_{CT} are allowed from QCD factorization, C/T up to 0.45 (0.55) for $\rho_A = 1(2)$. However the range of allowed values for C/T is still below the fitted values shown in Figure 1 for $\gamma > 47^\circ$. We thus obtain a constraint $\gamma \leq 44^\circ(52^\circ)$ and $\gamma \leq 46^\circ(56^\circ)$, for $\rho_A = 1$ and $\rho_A = 2$ respectively. In [20] the problem of the small a_2 has already been recognized and a possible solution was proposed : by choosing the the largest value of the Gegenbauer moment of π distribution amplitude




Figure 2: Scattered plot of the QCD factorization estimate for δ_{CT} (x -axis) versus C/T (y -axis) including the end-point singularity effects. In the plot, we fix the ρ parameters as $(\rho_H, \rho_A) = (1, 1)$ (left) and $= (1, 2)$ (middle) and vary the phases in the range of $-\pi < \phi_{A,H} < \pi$ (interval of 0.2 radian). The rest of the parameters are fixed (see text for details). The last figure (right) is obtained in the same manner with $(\rho_H, \rho_A) = (1, 1)$ but with different parameter set, the so-called scenario 2 of QCD factorization (see text for details).




Figure 3: The left figure is the plot of Eq. (34), the result including the FSI phase, in the plane of $\delta_{CT_{\text{eff}}}$ (x -axis) versus $C_{\text{eff}}/T_{\text{eff}}$ (y -axis) by varying δ_{20} . The number on the line indicates the value of δ_{20} at each point. The bare value $C_0/T_0 = 0.29$ obtained from the default parameter sets of the QCD factorization with $\rho_{H,A} = 0$ is used. In the middle figure, we put together the left figure and the experimental bounds from R_{+-} and R_{00} for the case of $\gamma = 57^\circ$ (Fig.1 upper-right). The right figure is obtained in the same way as the middle one but using $C_0/T_0 \simeq 0.61$, the result with the parameter set called scenario 2 in QCD factorization.

(scenario S2). More recently, this model has been reanalyzed with NLO corrections to the hard spectator-scattering diagram [21] and an enhancement by a factor of 2 for the hard-scattering correction is obtained:

$$a_2 \simeq 0.48e^{-i10^\circ} + 0.18\rho_H e^{i\phi_H} \quad (31)$$

$$C_0/T_0 e^{i\delta_{CT_0}} \simeq 0.61e^{-i3^\circ}, \rho_{H,A} = 0. \quad (32)$$

As shown in the scattered plot of Fig. 2 with $\rho_A = 1$ for scenario S2(right most), $C/T \simeq 1.1$ can be achieved in this scenario if δ_{CT} is very small. It appears that QCD factorisation can solve the large C/T puzzle.

4 Does FSI Phase Make C/T large

Recent studies show that FSI phase in $B \rightarrow \pi\pi$ can be large[22] and could effectively enhances C/T [23, 24], since the $B \rightarrow \pi^0\pi^0$ decay can be induced by the charge exchange scattering process $\pi^+\pi^- \rightarrow \pi^0\pi^0$ which can generate C from T . We examine here whether the FSI effect can enhance sufficiently the value of C/T of the QCD factorization without adjusting the incalculable parameters $\rho_{H,A}$ and $\phi_{H,A}$. We now add the FSI phases to the QCDF amplitude and evaluate C/T and constrain the values of $\delta_{20} = \delta_2 - \delta_0$ and δ_{CT} using the allowed region for C/T in Fig.1 with $\gamma = 57^\circ$. In [25], a similar analysis is performed and a large δ_{20} is found by a fit to the the central values of the experimental data. We have

$$\begin{aligned} T_{\text{eff}} e^{i\delta_{T_{\text{eff}}}} &= [(2T_0 - C_0)e^{i\delta_0} + (T_0 + C_0)e^{i\delta_2}]/3 \\ C_{\text{eff}} e^{i\delta_{C_{\text{eff}}}} &= [-(2T_0 - C_0)e^{i\delta_0} + 2(T_0 + C_0)e^{i\delta_2}]/3 \\ P_{\text{eff}} e^{i\delta_{P_{\text{eff}}}} &= P_0 e^{i(\delta_D + \delta_{0P})}. \end{aligned} \quad (33)$$

where C_0, T_0, P_0 are C, T and P for $\rho_{H,A} = 0$

$$\left(\frac{C_{\text{eff}}}{T_{\text{eff}}}\right) e^{i\delta_{CT_{\text{eff}}}} = \frac{(-2 + 2e^{i\delta_{20}}) + (1 + 2e^{i\delta_{20}})C_0/T_0}{(2 + e^{i\delta_{20}}) + (-1 + e^{i\delta_{20}})C_0/T_0} \quad (34)$$

$(C/T)_{\text{eff}}$ indeed becomes larger as the FSI phase increases. The bare ratio $C_0/T_0 = 0.29$ can be enhanced to $(C_{\text{eff}}/T_{\text{eff}}) \simeq 0.4$ for $\delta_{20} \simeq \pm 21^\circ$, where $\delta_{CT_{\text{eff}}} \simeq \pm 44^\circ$.

For $\gamma = 57^\circ$, $C_0/T_0 = 0.29$, the allowed region from R_{+-} and R_{00} overlap at $\delta_{20} \simeq -65^\circ$, at which $(C/T)_{\text{eff}} \simeq 0.96$ (middle figure). For $\gamma = 57^\circ$, $C_0/T_0 = 0.61$ (scenario S2), the central value of (R_{+-}, R_{00}) are reproduced by $\delta_{20} \simeq 40^\circ$ where $C_{\text{eff}}/T_{\text{eff}} \simeq 0.8$ and $\delta_{CT_{\text{eff}}} \simeq 40^\circ$. To constrain γ , we need the FSI phase and input from QCDF. Then assuming $\delta_{20} \lesssim 30^\circ$, we obtain $\gamma \lesssim 48^\circ (55^\circ)$ using the QCDF default values. With the parameters for scenario 2 shown above, we find that $\delta_{20} \lesssim 30^\circ$ leads to $\gamma = (59 \pm 3)^\circ (> 56^\circ)$. Note that there may be a FSI contribution not only to the phase but to C/T itself[24], so the bounds obtained above may receive corrections from QCD or FSI effects.

5 Conclusion

Using the latest $B \rightarrow \pi\pi$ data, we find that in $B \rightarrow \pi\pi$ decays C/T is large over a large range for δ_{CT} : $C/T \gtrsim 0.5$ for $\gamma > 47^\circ$. We then show that QCDF with large $\rho_{H,A}$ leads to large values of C/T , especially when δ_{CT} is small. In a scenario of QCDF with $\rho_{H,A} = 0$ but with the FSI strong phase included, C/T and δ_{CT} are enhanced when the FSI phase δ_{20} increases. A precise measurement of the $B^0 \rightarrow \pi^0\pi^0$ direct CP asymmetry C_{00} could distinguish these two models. In fact

$$\frac{C_{00}}{C_{+-}} = \frac{C}{T} \frac{\sin(\delta_{CT} - \delta_{PT})}{\sin \delta_{PT}} \frac{1}{R_{00}}. \quad (35)$$

A small $\delta_{CT} (\simeq 0)$ leads to a ratio of order unity with negative sign, $C_{00}/C_{+-} \simeq -(C/T)/R_{00}$, implying $C_{00} = 0.57$ for $\delta_{CT} = 0$, close to the higher end of the current experimental value $C_{00} = 0.28_{-0.39}^{+0.40}$ [19] while a large $\delta_{CT} (\simeq \pm\pi/2)$ would imply a strong dependence on δ_{PT} for the ratio $C_{00}/C_{+-} \simeq \pm(C/T)/(R_{00} \tan \delta_{PT})$.

6 Acknowledgments

I would like to thank Giulia Ricciardi and The Organizers for the Invitation to The Workshop and for the warm hospitality extended to me at Capri.

References

- [1] G. Buchalla, A. J. Buras, and M. E. Lautenbacher, *Rev. Mod. Phys.* **68** (1996)1125.
- [2] M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, *Phys. Rev. Lett.* **83** (1999) 1914 ; *Nucl. Phys. B* **591** (2000) 313 .
- [3] P. Ball, S. Khalil and E. Kou, *Phys. Rev. D* **69** (2004) 115011 [arXiv:hep-ph/0311361].
- [4] E. Kou and T. N. Pham, hep-ph/060113, to appear in *Physical Review D*.
- [5] M. Gronau and J. L. Rosner, *Phys. Rev. D* **65** (2002) 093012; *ibid* **66** (2002) 053003, [Erratum-*ibid.* *D* **66** (2002) 119901].
- [6] A. Ali, E. Lunghi and A. Y. Parkhomenko, *Eur. Phys. J. C* **36** (2004) 183 [arXiv:hep-ph/0403275].
- [7] Y. Y. Charng and H. n. Li, *Phys. Rev. D* **71** (2005) 014036.
- [8] P. Zenczykowski, *Phys. Lett. B* **590** (2004) 63.
- [9] C. W. Chiang, M. Gronau, J. L. Rosner and D. A. Suprun, *Phys. Rev. D* **70** (2004) 034020.
- [10] S. Mishima and T. Yoshikawa, *Phys. Rev. D* **70** (2004) 094024
- [11] X. G. He and B. H. J. McKellar, arXiv:hep-ph/0410098.
- [12] M. Gronau, E. Lunghi and D. Wyler, *Phys. Lett. B* **606** (2005) 95.
- [13] A. J. Buras, R. Fleischer, S. Recksiegel and F. Schwab, *Acta Phys. Polon. B* **36** (2005) 2015 [arXiv:hep-ph/0410407].
- [14] Z. z. Xing and H. Zhang, *Phys. Rev. D* **71** (2005) 051302.
- [15] M. Sowa and P. Zenczykowski, *Phys. Rev. D* **71** (2005) 114017.
- [16] Y. Grossman, A. Hocker, Z. Ligeti and D. Pirjol, arXiv:hep-ph/0506228.
- [17] M. Imbeault, arXiv:hep-ph/0505254.
- [18] G. Raz, arXiv:hep-ph/0509125.
- [19] Heavy Flavor Averaging Group, Winter 2005. arXiv:hep-ex/0505100.
- [20] M. Beneke, and M. Neubert, *Nucl. Phys. B* **675** (2003) 333.
- [21] M. Beneke and D. Yang, *Nucl. Phys. B* **736** (2006) 34; M. Beneke and S. Jager, arXiv:hep-ph/0512351.

- [22] M. Suzuki and L. Wolfenstein, Phys. Rev. D **60** (1999) 074019.
- [23] H. Y. Cheng, C. K. Chua and A. Soni, Phys. Rev. D **71** (2005) 014030; *ibid* **72** (2005) 014006.
- [24] S. Fajfer, T. N. Pham and A. Prapotnik Brdnik, Phys. Rev. D **72** (2005) 114001.
- [25] L. Wolfenstein and F. Wu, Phys. Rev. D **72** (2005) 077501.